

CONTRACEPTION–HORMONAL

COMBINED ORAL CONTRACEPTIVES (COCS)

The reason for practicing contraception (or birth control) is to prevent an unplanned pregnancy. The majority of methods of contraception enable sexually active couples to temporarily avoid pregnancy. Permanent birth control is accomplished through sterilization. There are many types of birth control methods that are currently available. There is no perfect method at the moment. Each has its own advantages and disadvantages. Be sure you know and understand the different types of birth control available to you, the risks and benefits of each, and any side effects, so that you can make an informed choice.

Combined oral contraceptives (COCs) pills prevent ovulation (release of a ripened egg from the ovary) and therefore make pregnancy unlikely or impossible. It also thickens the mucus in the cervix, making it difficult for sperm to reach an egg. The inner lining of the womb thins out during pills taking, so it is less likely to accept a fertilised egg.

The combination type, combining both oestrogen and a synthetic progestogen in a fixed daily dose, are the most popular and the most effective.

Effectiveness

The effectiveness is dependant on whether the pills are taken correctly, without missing a significant number of pills. If the pills are taken according to instructions, then it is over 99 per cent effective. This means that less than one woman in 100 using this method for a year will get pregnant.

Advantages

- Very effective, with lowest failure rate of any non-permanent method—less than 0.1 pregnancies per 100 women per year, if all pills are taken as prescribed.
- Periods become more regular.
- Less painful menstrual periods (reduction in period pain)
- Decrease in the amount of menstrual bleeding in most women.
- Decrease in incidence of cancer of ovary, womb and colon.
- May help with premenstrual symptoms
- Less likelihood of anaemia or arthritis.
- Decrease in incidence of pelvic inflammatory disease.
- Improvements in acne and oily skin.
- May reduce the risk of fibroids, ovarian cysts and (non-cancerous) breast disease

Disadvantages

- Not suitable for women who are smokers and above age of 35.
- Will not protect against sexually transmitted infections (STIs). A form of barrier protection will need to be used in those at risk.
- Not suitable for those who are forgetful to take the pills on a regular basis.
- May not be suitable for some women due to side effects.

Side effects

- There may be side effects from the pill. Many of these are not common and, even when they do, the frequency will decrease in severity after 1 to 2 months. Most patients experience no side effects at all.
- Common mild or minor side effect may include: headache, nausea, fluid retention, breast fullness or tenderness and mood changes. If these do not stop within a few months, changing the type of pill may help.
- Breakthrough bleeding (unexpected bleeding on pill taking days) and spotting is common in the first few months of pill use and will usually subside after a few cycles.
- Serious side effects are: stroke, heart attack and blood clots in the calf veins. Fortunately, these are very rare in healthy and non-smoking women.

Instructions – When and How to start?

There are many conditions that might make the pill unsuitable for you. Your doctor will take a thorough medical history and perform a physical examination prior to prescribing combined oral contraceptives. If any of the conditions are present that makes the pills unsuitable, another form of birth control will be recommended.

When?

- During menses – starts from day 1 onwards but not later than day 5.
- Starts immediately upon switching from other methods such as injectables type, implants or intrauterine device. You will be protected from pregnancy immediately.

How?

To start the first pack :

- Count the first sign of your menstrual bleeding as day 1.
- On either day 1, 2 or 3, begin taking 1 pill each day. Pick a time that suits you and stick to it as your routine. Use a reminder system or set an alarm (using your handphone/mobile device alarm) to help you remember. The pill can be taken without sips of water, if there is non available. The effect of pregnancy prevention by the pills is immediate. However, if you start the pill only after the 5th day, additional contraceptive protection (e.g using condoms) is required for

- 7 days before the pills is able to offer the pregnancy prevention effect.
- Continue taking the pills for 21 days.
 - Stop taking the pills for 7 days (this is called the pill free interval).
 - Begin taking pills again for the next cycle after these 7 days or rest, no matter when menstrual bleeding begins or ends.
 - If you do not have menses during the pill free interval, do not start the next pack. See your doctor first for further evaluation. This can occur in a small minority of women and can be normal. However, pregnancy needs to be excluded first.
 - If for whatever reasons you want to delay your period, you can continue to the next pack without stopping (no pill free interval). There will be no bleeding as long as the pills are taken.
 - If you forget to take the pill/pills, please refer to the patient information leaflet on missed COCs pills.

See your doctor immediately if there is:

- Increased frequency, amount, or duration of menstrual bleeding.
- Painful during intercourse or bleeding afterward.
- Pain and swelling in the calf of your leg, or any unusual leg pains.
- Severe chest pain or shortness of breath.
- Blood pressure goes up.
- Symptoms of gallbladder disease: upper abdominal pain, fever, jaundice, nausea.
- Any signs or symptoms that make you suspect pregnancy.
- See flashing lights or blurred images, or your ability to see decreases.
- Severe, throbbing headache.

Disclaimer

This is for informational purposes only and is not intended to be a substitute for professional medical advice, diagnosis, or treatment. It is important for readers to seek proper medical advice when necessary.

Dr Lee Say Fatt
Sime Darby Medical Centre, Subang Jaya
Revised July 2011